



## Uses of Difenoconazole 25% EC

Crop	Common name of the disease	Dosage per ha			Waiting period from last application to harvest (in days)
		a. i. (g)	Formulation (g/ml)/%	Dilution in water (L)	
<b>Difenoconazole 25% EC</b>					
Apple	Scab	0.004% or 4 g/100 ltr. water	0.015% or 15ml/100ltr. water	As required depending upon the size of the plant and plant protection equipment used	14
Paddy (Rice)	Sheath blight	0.0125% or 12.5g/100 ltr. water	0.05% or 50 ml/100 ltr. water	500	25
Chilli	Die-back, Fruit rot	0.0125% or 12.5g/100 ltr. water	0.05% or 50 ml/100 ltr. water	500-1000 (as per the size of plant )	15
Cumin	Blight, Powdery mildew	0.0125% or 12.5 g/100 ltr.. water	0.05% or 50 ml/100 ltr..water.	500	15
Onion	Purple blotch	0.025% or 25 g/100 ltr. water	0.1% or 100 ml/100 ltr..water.	500	20
Pomegranate	Fruit rot	0.025% or 25g/100ltr. water	0.1% or 100 ml/100 ltr. water.	500	7
Grape	Anthracoese, Powdery mildew	0.0075% or 7.5 g/100 ltr. water	0.03% or 30ml/100 ltr. of water	500	10