



Uses of Metalaxyl 35% WS

Crop	Common name of the disease	Dosage per ha			Waiting period from last application to harvest (in days)
		a. i. (g)	Formulation (g/ml)/%	Dilution in water (L)	
Metalaxyl 35% WS					
Maize	Sorghum Downy mildew, Sugarcane downy mildew, Phillippine Downy mildew, Brown stripe downy mildew	Slurry seed treatment with 240g/100 kg seed	700g/100 Kg seed	0.75-1.0/100 Kg seed	3½-4 months Depending on the variety
Bajra	Downy mildew	Slurry seed treatment with 200g/100 Kg seed	600g/100 Kg seed	0.75-1.0/100kg seed	3-3½ months Depending on the variety
Sorghum	Downy mildew	Slurry seed treatment with 200g/100 Kg seed	600g/100 Kg seed	0.75-1.0/100kg seed	3½-4months Depending on the variety
Sunflower	Downy mildew	Slurry seed treatment with 200g/100 Kg seed	600g/100 Kg seed	0.75-1.0/100kg seed	3½-4months Depending on the variety
Mustard	White rust	Slurry seed treatment with 200g/100 Kg seed	600gm/100 Kg seed	0.75-1.0/100kg seed	3½-4months Depending on the variety