Uses of Propineb 70% WP

Crop	Common name of the disease	Dosage per ha			Waiting
·		a. i. (g)	Formulation (g/ml)/%	Dilution in water (L)	period from last application to harvest (in days)
Propineb 70	% WP				
Apple	Scab	0.21% or 210 g/100 Ltr. water	0.30% or 300 gram/100 Ltr.water	As required depending upon size of the tree and plant protection equipment used	30
Pomegranate	Leaf and Fruit spots	0.21% or 210 g/100 Ltr. water	0.30% or 300 gram/100 Ltr.water	As required depending upon size of the tree and plant protection equipment used	10
Potato	Early & Late blight	0.21% or 210g/100 Ltr. water	0.30% or 300 gram/100 Ltr.water	As required depending upon crop stage and plant protection equipment used	15
Chilli	Die-back	0.21% or 210 g/100 Ltr. water	0.5% or 500 gram/100 Ltr.water	As required depending upon crop stage and plant protection equipment used	10
Tomato	Buck eye rot	0.21% or 210 g/100 Ltr. water	0.30% or 300 gram/100 Ltr.water	As required depending upon crop stage and plant protection equipment used	10



PERFECT SOLUTIONS

Office Add.: B-371, Paschim Vihar, New Delhi - 110063 (India) Factory Add.: 15/1, 15/2, 15/3 Industrial Estate, Vidisha - 464001 (M.P.)

Email: info@perfectsolution.best

Grapes	Downy mildew	0.21% or	0.30% or 300	As	40
		210 g/100	gram/100	required	
		Ltr. water	Ltr.water	depending upon	
				crop stage and	
				plant protection	
				equipment used	