



Uses of Fenoxaprop-p-ethyl 9.3% EC

Agricultural Use

Herbicide name & approved Crops	Weed species	Dosage/ha			Waitig period/ PHI between last application & harvet (days)
		a.i. (gm/ Kg)	Formulation in (gm/ml /Kg/ ltr)	Dilution in Water (Liter)	
Fenoxaprop-p-ethyl 9.3% EC					
Soybean	<i>Echinochloa colonum</i> , <i>Echinochloa crusgalli</i> , <i>Digitaria sp.</i> , <i>Eleusine indica</i> , <i>Setaria sp.</i> , <i>Brachiaria sp.</i>	100gm.	1111 ml. (15-20 DAS)	250-300	100
Rice (enticulate)	<i>Echinochloa crusgalli</i> , <i>Echinochloa colona</i>	56.25 gm	625 ml. (10-15 DAT)	300-375	70
Blackgram	<i>Echinochloa crusgalli</i> , <i>Echinochloa colona</i> <i>Digitaria sp.</i> <i>Dactyloctenium</i> <i>Aegyptium</i>	56.25-67.5 g	625-750ml. (15-20 DAS)	375-500	43
Cotton	<i>Echinochloa sp.</i> <i>Eluesine indica</i> <i>Dactyloctenium</i> <i>Aegyptium</i> <i>Eragrostit minor</i>	67.5 g	750ml. (20 -25 DAS)	375-500	87
Onion	<i>Echinochloa colonum</i> <i>Dactyloctenium aegyptium</i>	78.75	875	375	10
Groundnut	<i>Echinochloa sp.</i>	78.75	875	375	89