



Uses of Metribuzin 70% WP

Agricultural Use					
Herbicide name& approved Crops	Weed species	Dosage/ha			Waitig period/ PHI between last application & harvet (days)
		a.i. (gm/ Kg)	Formulati on in (gm/ml /Kg/ ltr)	Dilution in Water (Liter)	
Metribuzin 70% WP					
Soybean	<i>Digitaria spp.</i> <i>Cyperus esculentus</i> <i>Cyperus campestris</i> <i>Borreria spp.</i> <i>Eragrostis spp.</i>	0.35-0.525 kg	0.5-0.75kg.	750-1000	30
Wheat	<i>Phalaris minor</i> <i>Chenopodium album</i> <i>Melilotus spp.</i>	Medium soil-0.175kg Heavy soil - 0.21kg	0.25 kg 0.30 kg.	500-750	120
Sugarcane	<i>Cyperus rotundus</i> , <i>Cynodon dactylon</i> , <i>Astodelus fistulosus</i> , <i>Chenopodium album</i> , <i>Convolvulus arvensis</i> , <i>Portulaca oleracea</i> , <i>Anabalis servensis</i> , <i>C.intybus</i> , <i>Echinochloa colonum</i> , <i>Dactyloctenium aegyptium</i> , <i>Parthenium hysterophorus</i> , <i>Commelinia spp.</i>	Post-emergence, 25-30 DAP Or Pre-emergence At first workable spray condition or 3-5 DAP	1.05-1.4 1.05-2	1.5-2 1.5-3	60



PERFECT SOLUTIONS

Office Add.: B-371, Paschim Vihar, New Delhi - 110063 (India)

Factory Add.: 15/1, 15/2, 15/3 Industrial Estate, Vidisha - 464001 (M.P.)

Email: info@perfectsolution.best

Potato	<i>Chenopodium album,</i> <i>Trianthema monogyna,</i> <i>Parthenium hysterophorus,</i> <i>Funaria parviflora,</i> <i>Melilotus spp., Phalaris minor</i>	Pre-emergence 3-4 DAP or Post-emergence between emergence and time when the first potato plants have reached height of 5 cms.	0.525	0.750	30
Tomato	<i>Trianthema portacastrum,</i> <i>Dactylectonium aegypticum,</i> <i>Gynandropsis pentaphylla,</i> <i>Amaranthus viridis,</i> <i>Portulaca oleracea,</i> <i>Digera arvensis,</i> <i>Euphorbia frustucula</i>	Pre-planting a week before transplanting Or	0.525	0.760	30